

A Study on the Feasibility and Utility of Continuous Glucose Monitors in Elite Football

Sophie Harries^{1,*}, David Clayton¹, Michael Johnson¹, Ross Burbeary², Robin Thorpe³, Ian Varley¹

¹School of Science and Technology, Nottingham Trent University, Nottingham, Clifton Campus, Clifton Ln, Clifton, Nottingham NG11 8NS

²Huddersfield Town Association Football Club, Millers Oils High Performance Complex, 509 Leeds Rd, Huddersfield HD2 1YJ

³Red Bull Performance Centre, 2700 Pennsylvania Ave, Santa Monica, CA 90404, United States

Research Article

Open Access &

Peer-Reviewed Article

DOI: 10.14302/issn.2694-2283.jsem-24-5363

Corresponding author:

Sophie Harries, School of Science and Technology, Nottingham Trent University, Nottingham, Clifton Campus, Clifton Ln, Clifton, Nottingham NG11 8NS

Keywords:

Glucose, Performance, Continuous Glucose Monitoring, Football, Soccer

Received: November 28, 2024

Accepted: December 26, 2024

Published: January 16, 2025

Citation:

Sophie Harries, David Clayton, Michael Johnson, Ross Burbeary, Robin Thorpe et al. (2025) A Study on the Feasibility and Utility of Continuous Glucose Monitors in Elite Football. Journal of Sports and Exercise Medicine - 1(2):8-21. <https://doi.org/10.14302/issn.2694-2283.jsem-24-5363>

Abstract

Physiological performance may be limited by reduced systemic glucose availability in working muscles. Continuous glucose monitors (CGM) measure interstitial glucose every 1- 15 minutes, offering a practical way to assess glucose during sporting activity. However, empirical research has predominantly focused on endurance-based sports, glucose responses during a professional competitive football match remain unknown. This study evaluated the feasibility and utility of CGM feasibility in professional football. Eight professional, male outfield footballers from the English third tier participated in the study. Participants completed a 14-day food diary control period, followed by a 28-day observation period wearing CGM devices during six matches and sixteen training sessions. CGM devices remained *in situ* for 98% of training sessions and matches. Mean glucose concentrations were 6.5 ± 1.2 mmol/L during training sessions, 7.5 ± 2.1 mmol/L during match play, and 5.4 ± 0.3 mmol/L overnight. No significant differences were found between glucose concentrations during match-play ($P = 0.060$) or training ($P = 0.510$), compared to overnight fasted glucose concentrations. There was also no difference between training and match-play glucose concentrations ($P = 0.788$). Glucose concentrations were highly individualised, with one player displaying minimal change throughout match play (-0.2 mmol/L) whereas another experienced increases of 5.8 mmol/L. Non-nutritional factors appeared to influence glucose levels; for example, participants ($n=3$) who used nicotine pouches displayed an acute increase in blood glucose in the 10-60 minutes after administration. This study concludes that CGM use in professional football is feasible and valuable for assessing individual glucose responses to training and match-play.

Introduction

Intermittent team sports and endurance sports, rely on glycogenolysis to maintain blood glucose concentrations [1]. Sustained periods of muscle glycogenolysis

lead to significant muscle glycogen depletion (Krustrup et al., 2011), progressively reducing energy substrate availability in fast-twitch muscle fibres critical for explosive movements. This metabolic exhaustion directly correlates with peripheral muscle fatigue development (Knuiman et al., 2015), potentially compromising player performance by limiting power output and increasing perceived exertion in later match stages (Hudson et al., 2021). Research by Krustrup et al. (2011) empirically documents these metabolic changes, demonstrating that muscle glycogen levels can decrease by 40-60% during a typical match, depending on playing intensity and individual metabolic characteristics. Reduced glucose availability, as a result of either low blood glucose, low glycogen stores or low carbohydrate intake, is also associated with the development of central fatigue [1] due to reduced brain glucose uptake [2]. Preserving blood glucose concentration can reduce the reliance on glycogen for carbohydrate oxidation, and may therefore delay an aspect of fatigue during match-play, benefitting tactical and physical performance [3]. Although the relationship between blood glucose concentrations and muscle glycogen is not fully understood [4], maintaining blood glucose increases carbohydrate availability, and helps to meet the metabolic demand of exercise [5]. However, the optimal blood glucose concentration for exercise performance is not known and therefore the monitoring glucose concentrations in a sporting population may be of value.

Traditionally, blood glucose concentrations have been measured using fingertip capillary blood sampling. In most contexts, this is impractical, causes discomfort, and only provides discrete data on a measure that changes continually during exercise [6]. Continuous Glucose Monitors (CGM) are now routinely used within healthcare services to provide a continuous measurement of interstitial glucose concentrations. This has revolutionised healthcare for people with diabetes [7], with real-time interstitial glucose data helping people to monitor glucose concentrations and patients to avoid hyper (>10 mmol/L) or hypoglycaemia (<3.9 mmol/L) without medical intervention [8]. In the context of sport, the ability to continuously monitor glucose concentrations pre, during and post-exercise may help to inform fuelling strategies related to carbohydrate type and timing of ingestions with the aim of optimising performance and recovery (Holzer et al., 2022). However, the benefits of using CGM in a sporting setting is far from established.

Research using CGMs in professional sport is emerging. Previous studies evaluating CGM have focussed on sports where a need for high carbohydrate availability is most well established in relation to performance, such as endurance cycling [9] and running [10]. Studies employing CGMs have reported high levels of individuality in glucose concentrations, beyond that of measurement errors, in response to both exercise and carbohydrate intake even with a standardised diet [9, 10, 11]. Individuality in response could be influenced by stress, illness, dehydration or hormones [12], therefore identifying individualised dietary or exercise responses can be hindered by non-dietary factors that influence glycaemic control. Carbohydrate absorption, metabolism, uptake and utilisation (Guest et al., 2019), eating behaviour [13], and commonly reported differences in gut microbiome [14] could also be impacted by genetic variation. Therefore, it could be argued that biometric feedback [15] could be used for athletes to tailor a nutrition plan that accounts for individual differences. Previous studies have reported limited adverse effects of CGM use, and the data from CGM have been used to inform athlete fuelling and race pacing strategies [9, 10]. Based on these findings, CGM may be used within endurance sports to allow for bespoke fuelling strategies that help to mitigate fatigue and improve athletic performance. Of note, the use of CGMs during competitions have recently been banned in professional cycling due to concerns around personal privacy, financial accessibility, and to reduce the influence of technology on human racing [31]. While these factors should be carefully considered by

governing bodies, the existing research demonstrates the potential for biometric information to positively inform bespoke fuelling strategies within endurance sport. Other sports that involve also involve an endurance component, such as team sports, could similarly benefit from biometric feedback devices. Unfortunately, there is limited research assessing their practical use. For example, the research conducted in football is restricted to simulated match-play of semi-professional players using a standardised treadmill protocol (Hiromatsu et al., 2023). However, this limits the extent to which the data collected can be generalised to a competitive match because these methods do not account for the contact nature of competitive football (which could dislodge sensors), and within-player variability in metabolic demand due to the variable dynamics of professional match-play [16]. Therefore, the aim of the present study was to evaluate the feasibility and utility of CGM device use within professional footballers. Feasibility was defined as retention of CGM devices during an in-season competitive period. Utility was defined as the potential for the biometric data collected to impact nutrition strategies. A secondary aim was to determine the interstitial glucose concentrations during training and match play.

Methods

Participants

Eight professional male (age: 28.0 ± 5.5 years, body mass 86.4 ± 4.8 kg, height: 1.8 ± 0.1 m) outfield footballers competing in the third tier of English football during the 23/24 season volunteered for the study. Participants had no history of diabetes and provided written informed consent. Ethical approval was granted by Nottingham Trent University Ethics Committee.

Study Design and Procedures

Eight participants completed a 14-day control (112 total player days) period, during which they reported dietary intake. This was followed by a 28-day observation period (188 total player days), whereby participants wore the CGM device (Abbot Libre Sense Glucose Sport Biosensor, United States) and continued to report dietary intake. Two participants withdrew from the study during the observation period (Figure 1).

Continuous Glucose Monitors

CGM devices are reported to have a lag time of 2.4 ± 4.6 minutes and >90% of sensor readings are within 20% of blood glucose concentrations [17]. The device has an upper limit of 11.2 mmol/L. The device was inserted into the back of the upper arm by the researcher and replaced after 14 days, as per manufacturer instructions [18]. Specialist manufacturer-supplied performance patches, made of a silicone protection ring and 4-way stretch kinesiology tape, were placed onto the devices for protection. The CGM device was paired with the user's smartphone application [31] via Bluetooth for minute-by-minute data collection. If the device was disconnected from Bluetooth (e.g., due to distance between participants and their smartphone), readings were taken every 15 minutes and stored by the device for 8 hours until it was re-synchronised via Bluetooth. Post-training, at half-time and post-match, participants were instructed to re-synchronise the CGM device with their smartphone via Bluetooth.

Dietary logs

Participants reported dietary intake using a Snap and Send method, aimed to reduce reporting bias [19]. Participants took photos of all food and drink consumed alongside an object of known size (massage ball, 8cm diameter) at 45- and 90- degree angles. Participants received individual demonstrations regarding the dietary reporting method with common troubleshooting issues discussed, e.g., complex




Figure 1. Participant flow diagram

dishes with multiple ingredients to be listed alongside the photograph. Images were sent via an instant messenger service [21] to the primary researcher. If participants had not sent a photo through within 2 hours of their usual predicted mealtime, the researcher reminded participants via a text message e.g., breakfast reminder was sent at 10:30 in anticipation that a meal would have been consumed around 08:30. If participants failed to photograph their meal, dietary recall information was provided with portion sizes estimated. The primary researcher was present when meals were consumed at the training centre and assisted with dietary reporting by photographing and documenting supplement use. Participants reported all supplement use as part of their food diary, examples include caffeine gum, beta-alanine, creatine, and electrolytes. If supplements were provided for the participants by the club, dosage was confirmed with the team's nutritionist. For supplements taken at home, participants reported brand, weighed measure or dosage. On Day 2 of the observation period, the potential impact of tobacco free nicotine pouches, alongside tobacco containing Snus, on glucose concentration was observed and, therefore, nicotine pouch/snus use was reported on Day 2 onwards. Nicotine pouch/snus administration times and corresponding CGM value were time aligned based on the known delay (5 minutes) with CGM readings [17].

Training Load

During the observation period, three home matches and three away matches were played. Training schedules (16 sessions) and match day squad selection were not adjusted or influenced based on CGM study involvement or findings. Global Positioning System (GPS) units (Vector, Catapult, Australia) were positioned between the shoulder blades using a manufacturer-supplied vest and worn throughout training and match-play. Post-training and match-play, each GPS data set was downloaded and analysed using commercially available software (Viper, STATSports, Ireland). The physical performance variables assessed included: total distance covered (m), high speed (>5.5 m/s) distance covered (m), sprint speed (>7.0 m/s) distance covered (m), number of accelerations above 0.5 m/s² for >0.5 s, and number of decelerations below -0.5 m/s² for >0.5 s.

Feasibility

The feasibility of CGM devices was assessed based on the percentage of completed training and match sessions during which the CGM device remained in situ with data available for analysis. Additionally, any adverse effects reported by participants, particularly discomfort, were recorded.

Statistics

Overnight fasted glucose concentrations were determined by calculating the mean of overnight glucose concentrations from 00:00-06:00 during Week 1 of the observation period. This mean was used as a baseline measure. Glucose concentrations were assessed three hours pre-match, first half, half-time, second half, three hours post-match, and for training sessions, for each participant. Mean concentrations are reported for each period. During training and match play, the Bluetooth connection to CGM was lost, due to the distance between the monitor and the player's smartphone, so 1 5-minute mean glucose concentrations were recorded. As such for match play, first and second half match play was restricted to three measurements in 15-minute intervals. Percentage and absolute change from overnight fasted interstitial glucose concentrations were calculated for all match-play and training session measurements.

A Shapiro-Wilk's test was conducted to assess normality in distribution of data with all data deemed as normally distributed. Paired T-tests, with a significance level of $p < 0.05$, and Bonferroni correction were used to assess the difference between mean glucose concentrations during training sessions and match play (inclusive of 1st half, half time and 2nd half), overnight fasted glucose concentrations and training, and overnight fasted glucose concentrations and match-play. A paired T-Test was also used to assess any difference in dietary intake between the control and observation period.

Dietary intake was analysed by a Sports and Exercise Nutrition registered (SENr) dietitian, using dietary analysis software [30]. Dietary intake was quantified as energy (kilocalories), carbohydrates (grams) and protein (grams) in absolute units and relative to body mass. Mean energy and macronutrient intake between the control and observation period were compared to establish any significant changes for the duration of the study using a paired T-test, following normality testing [20].

Results

Feasibility

CGM were used in 148 training or match events, and needle displacement occurred on three occasions, all during training, thus the CGM produced data for 98% of training and match-play events without displacement. Six participants regularly participated in training and match-play. Four of the six participants sustained short-term muscular strain injuries, which resulted in a total of 9 player days of modified training i.e 'off feet' indoor gym work. Two participants were injured prior to the study start

Table 1. Dietary intake values for energy, carbohydrates and protein across the study duration (Overall), Match Day-1, Match Day and Match Day+1. Values are mean \pm SD and expressed in absolute and relative units (n=8)

	<i>Overall</i>	<i>Match Day-1</i>	<i>Match Day</i>	<i>Match Day +1</i>
Energy (kcal/day)	2704 \pm 410	3377 \pm 644	2808 \pm 759	2024 \pm 387
Energy (kcal/kg BM)	31.3 \pm 4.7	39.0 \pm 6.6	32.7 \pm 9.5	23.5 \pm 4.9
Carbohydrates (g/day)	270 \pm 53	369 \pm 96	317 \pm 71	176 \pm 38
Carbohydrates (g/kg BM)	3.1 \pm 0.6	4.2 \pm 0.9	4.2 \pm 0.9	2.1 \pm 0.6
Protein (g/day)	161 \pm 34	201 \pm 46	146 \pm 6	115 \pm 27
Protein (g/kg BM)	1.9 \pm 0.4	2.3 \pm 0.5	1.7 \pm 0.7	1.3 \pm 0.3

date and did not return to training or match-play during the study period, participating only in rehabilitation sessions. However, these participants continued to wear the CGM and record dietary intake throughout their rehabilitation.

Dietary Intake

Carbohydrate intake did not differ between the control and observation periods (control 3.0 \pm 0.6 g/kg/day, observation period 3.2 \pm 0.7 g/kg/day) ($t_7 = 1.595$, $p = 0.077$). There was no difference in energy and protein intake between the control and observation periods ($t = 0.028$, $p = 0.489$ and $t = 0.016$, $p = 0.494$, respectively). Dietary intake values can be seen in Table 1. Dietary recall rather than the snap and send method was used for one participant two days preceding drop out.

Response to Training and Match Play

The demands of training (mean total distance = 3930 \pm 432m, mean accelerations = 59 \pm 13, mean decelerations = 47 \pm 10, mean high-speed running = 143 \pm 94m) and match-play (mean total distance = 6044 \pm 3700m, mean accelerations = 84 \pm 35, mean decelerations = 82 \pm 33, mean high-speed running = 294 \pm 256m) were captured during all sessions.

Mean glucose concentrations were 6.5 \pm 1.2 mmol/L during training sessions, 7.5 \pm 2.1 mmol/L during match play, and 5.4 \pm 0.3 mmol/L overnight. There were no significant differences in glucose concentrations in response to match-play ($p = 0.060$) or training ($p = 0.510$), compared overnight fasted interstitial glucose concentrations. There was also no significant difference between training and match-play glucose concentrations ($p = 0.788$) (Figure 2).

Inter-Participant Variability

Glucose concentrations during match-play varied between participants (Figure 2 and Table 2). During the first half of match-play the absolute change in glucose concentrations from overnight fasted glucose concentrations was between - 0.2 mmol/L and + 5.8 mmol/L and similar changes were seen during the second half of match play (- 0.2 mmol/L and + 5.8 mmol/L).

Effects of nicotine pouch use on glucose concentration

Participants who administered nicotine pouches and Snus (n=3) resulted in a mean 0.7 \pm 0.4 mmol/L




Figure 2. Box and Scatter plot with individual player means for blood glucose 3 hours Pre- match, 1st half match play, half-time, 2nd half match play and 3 hours post-match.

* Participant 2 (∇) reached the sensor threshold of 11.2 mmol/L

Table 2. Absolute and Percentage change in mean glucose concentrations during match-play compared to mean overnight fasted glucose concentrations.

Player ID	<i>1st half absolute change (mmol/L)</i>	<i>1st half percentage change</i>	<i>Half time absolute change (mmol/L)</i>	<i>Half-time percentage change</i>	<i>2nd half absolute change (mmol/L)</i>	<i>2nd half percentage change</i>
1	2.2	42.4	2.9	56.8	2.4	47.1
2	5.8	106.8	5.8	106.8	5.8	106.8
3	3	50.1	1	17.7	1.6	26.6
4	1.3	23.1	1.8	32.5	1.7	30.4
5	0.5	9.1	0.8	13.9	0.7	11.6
6	-0.2	-4.2	0	0.1	-0.2	-3.7

* Participant 2 reached the sensor threshold of 11.2 mmol/L




Figure 3. Glucose concentration before and after nicotine pouch administration (0 minutes) without the presence of diet or exercise factors.

($13.0 \pm 8.2\%$) increase in glucose concentrations compared to the overnight fasted concentration. Three singular changes in the glucose response are demonstrated in Figure 3. These changes occurred in the absence of dietary intake and/or physical activity at the time of administration.

Discussion

Feasibility

The primary aim of this study was to assess if the use of CGM technology was feasible within professional football. This study demonstrated that CGM devices can be feasibly used to monitor glucose concentrations in professional football, evidenced by successful recording in 98% of the training or match play events (135 hours and 57 minutes of match play and on-pitch training sessions). Previous studies have reported limited adverse effects of CGM use, and the biometric data from CGM devices used to inform athlete fuelling and race pacing strategies [9, 10]. Based on these findings, CGM may be used within sports to allow for bespoke fuelling strategies that optimise carbohydrate utilisation to help mitigate fatigue and improve athletic performance, through immediate insights into an athlete's physiological energy utilisation and response. In support of this, CGMs have recently been banned in professional cycling, due to concerns around personal privacy, financial accessibility, and to reduce the influence of technology on human racing [31]. These factors will need to be considered by all sporting governing bodies, but this does indicate a potential for CGM to positively influence athletic performance. CGM also revealed some individual variability in glucose concentrations during match play and some unexpected changes in glucose concentration in response to nicotine pouch administration.

In summary, these findings suggest there may be value of using CGM in professional footballers, to better understand the individual responses to nutrition and lifestyle and identify players who may benefit from personalised support strategies.

Response to Match Play

CGM identified individual differences in blood glucose response to training and match-play. The difference between the highest and lowest glucose concentrations was 5.9 mmol/L. The metabolic demands of football vary from match to match and between players [22] and this may partially

contribute to the individuality observed in the present study. However, individuality has also been reported in endurance athletes [9, 10], where metabolic demands may be more consistent. As observed by [28], day-to-day variability between athletes was more prominent than overnight variability, and secondary to diet and activity induced changes in glucose concentration. Variability between individuals is likely to be associated with genetic differences impacting carbohydrate absorption, metabolism, uptake, and utilisation (Guest et al., 2019). Based on the variability observed in the present and previous studies, monitoring individual glucose responses using a CGM device could allow practitioners to develop personal fuelling strategies that account for individual variation in glucose availability during exercise (Hulton et al., 2022; Russell & Kingsley, 2014). Despite this, there is no existing data on 'optimal' blood glucose range for performance in football, so the value of monitoring and/or responding to glucose changes is debatable. Hiromatsu et al., (2023) concluded that it is likely important to maintain high glucose availability during match-play, to ensure provision of a continuous source of glucose to fuel carbohydrate metabolism, required for high-intensity efforts [1]. Therefore, the value of monitoring glucose concentrations via CGM for performance in football remains to be established.

CGM has improved care for people living with diabetes, with multiple studies demonstrating improvements in glycaemic control and reduced reliance on glucose-lowering medication in patients using CGM [7]. Much of this success is attributed to the educational value these monitors provide patients, encouraging changes in diet and physical activity to improve glucose concentrations, leading to improved health outcomes [23]. Whether continuous monitoring of glucose concentrations via CGM can influence behaviour in athletes is currently unknown. Many athletes struggle to achieve the carbohydrate intake recommendations [24], with our study finding that players did not reach the UEFA recommended carbohydrate intake of 6-8 g/kg BM [25]. Interestingly we did observe a non-statistically significant tendency ($p = 0.077$) for carbohydrate to increase slightly during the CGM observation period.

Although the absolute difference is small (0.2 g/kg BM), there are known limitations of food diaries to accurately capture intake [18], and as such, this finding could be interpreted as indicative of change rather than an accurate reflection of absolute intake.

Further research is required in a larger sample size and over a longer time frame to better understand the potential educational value of CGM in athletes to positively influence dietary intake behaviours.

One significant consideration for practitioners surrounding the utility of CGM devices is that during match-play CGMs are not synchronised. This results in a mean reading being recorded every 15 minutes rather than minute by minute. There is also an approximate 5-10 minute delay before interstitial glucose concentrations reflect blood glucose concentrations [16], which needs to be accounted for by practitioners. However, with sufficient duration and repetition, there may be potential for CGM to generate a 'blood glucose profile' for individual players, which could allow for proactive nutritional approaches.

Other Insights

An advantage of CGM is the capturing of unexpected effects of diet or behaviour on glucose concentrations. In the present study, three of the eight participants regularly used either a tobacco free nicotine pouch or tobacco containing snus pouch, administered between the gum and upper lip. Nicotine pouches and Snus are reported to help relaxation and reduce stress [26]. A recent report found 1 in 5 professional football players within the English Football league currently use nicotine pouches or

snus [26]. We observed that sublabial administration of the nicotine pouch/snus led to a 13% increase on mean in glucose concentrations up to 55 minutes post-administration. However, peak change was variable for each administration and participant as displayed by Figure 3. Nicotine increases lipolysis with theories surrounding this response is potentially driven through the inhibition of glucose uptake [27, 28].

Current recommendations emphasise the importance of rapid glycogen resynthesis within the first four hours post-match, with inadequate glycogen resynthesis leading to a reduction in performance for the next 48 - 72 hours [25]. As such, the novel findings from this study appear to suggest a high contributing role of nicotine interfering with glucose regulation, which could influence both glucose stability and glycogen resynthesis, with the potential for negative effects on recovery. Given the widespread use of nicotine pouches and snus in both professional and amateur football [26], further research is required to determine the effects of nicotine on blood glucose in a larger population and under controlled conditions.

Practical Considerations

Although CGM successfully recorded blood glucose on 98% of occasions, careful consideration is required on whether the performance insights gained outweigh the device cost and analysis time. However, there is value in discovering potentially unexpected changes in glucose concentrations not linked to diet or exercise, such as those seen in the current study with nicotine pouch use. Such observations may extend to pre- and post-match individualised carbohydrate recommendations and periodisation. These observations increase awareness and may enable support staff to implement preventative strategies.

CGM may increase player awareness of how diet and lifestyle practices influence their glucose concentrations, which could be used as a tool to encourage positive change. In the present study, carbohydrate intake in match day -1, was observed to be considerably lower than UEFA recommendations (observed 4.2 ± 0.9 g/kg BM, UEFA guidance 6-8 g/kg) [25], which is consistent with previous studies in football players (Danielik et al., 2022; Jenner et al., 2019). The present study observed a modest increase in CHO intake during CGM of 0.2 g/kg BM or 20 g per day. However, during statistical testing this increase to be non-significant. Despite this argument could be made that within the applied setting this tentatively supports CGM use to positively change lifestyle behaviours, as has been previously shown in clinical populations [23].

Some issues that may impact feasibility were identified. CGM devices are currently targeted for clinical use and include a built-in upper limit to prevent self-diagnoses of diabetes. In the present study, one of the six participants regularly reached/exceeded the maximal reading of 11.2 mmol/L. This is a high value that would not be expected to be regularly exceeded in a professional sport context. The inability to conduct minute-by-minute analysis during training and match-play, caused by infrequent Bluetooth connection between the CGM device and the user's smartphone, significantly limits the depth of data interpretation. It is still unclear whether glucose measurements taken every 15 minutes in these dynamic environments provide valuable insights for practitioners, given the rapid fluctuations in glucose levels. More consistent, real-time monitoring may be required to fully understand the physiological changes and provide meaningful guidance for performance and health management.

Conclusion

This study found that CGM devices can be feasibly used to monitor glucose concentrations in

professional footballers. As with other sports, individuality in glucose response was observed between participants, indicating that the insight provided by CGM could be valuable for informing personalised fuelling and recovery strategies. There may also be educational value to CGM, both to enhance players understanding of the relationship between dietary intake, glucose availability and performance, and for practitioners to identify practices that may negatively affect glucose regulation (e.g., nicotine administration).

Acknowledgments

The authors would like to thank the participants of this study for their time, effort, and co- operation of the participants of this study.

Author contribution (according to the CRediT):

Conceptualization – Dr Ian Varley, Dr David Clayton, Ross Burbeary. *Data curation* – Sophie Harries

Formal analysis – Sophie Harries

Funding acquisition – Dr Ian Varley, Dr David Clayton, Ross Burbeary *Investigation* – Sophie Harries

Methodology – Sophie Harries, Dr Ian Varley, Dr David Clayton, Ross Burbeary *Resources* – Dr Ian Varley, Dr David Clayton, Dr Michael Johnson, Ross Burbeary *Supervision* – Dr Ian Varley, Dr David Clayton, Dr Michael Johnson, Dr Robin Thorpe *Validation* – Sophie Harries, Dr Ian Varley, Dr David Clayton, Dr Michael Johnson *Visualization* – Sophie Harries, Dr Ian Varley, Dr David Clayton, Dr Michael Johnson

Writing – original draft – Sophie Harries, Dr Ian Varley, Dr David Clayton, Dr Michael Johnson

Writing – review & editing – Sophie Harries, Dr Ian Varley, Dr David Clayton, Dr Michael Johnson, Ross Burbeary, Dr Robin Thorpe.

Funding details

Funding for this project was provided by Nottingham Trent University as part of a PhD studentship.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Data availability statement

The data that support the findings of this study are openly available in Zenodo at <https://doi.org/10.5281/zenodo.14202932>

Code availability statement

The code that support the statistical analysis of this study are openly available in Zenodo at <https://doi.org/10.5281/zenodo.14202932>.

Ethics approval and Informed Consent

Ethics approval was granted from Nottingham Trent University (1849189).

References

1. Hargreaves, M., & Spriet, L. L. (2020). Skeletal muscle energy metabolism during exercise. *Nature Metabolism*, 2(9), Article 9. <https://doi.org/10.1038/s42255-020-0251-4>
2. Khong, T. K., Selvanayagam, V. S., Sidhu, S. K., & Yusof, A. (2017). Role of carbohydrate in

- central fatigue: A systematic review. *Scandinavian Journal of Medicine & Science in Sports*, 27(4), 376–384. <https://doi.org/10.1111/sms.12754>
3. Flockhart, M., & Larsen, F. J. (2023). Continuous Glucose Monitoring in Endurance Athletes: Interpretation and Relevance of Measurements for Improving Performance and Health. *Sports Medicine*. <https://doi.org/10.1007/s40279-023-01910-4>
 4. Sylow, L., Kleinert, M., Richter, E. A., & Jensen, T. E. (2017). Exercise-stimulated glucose uptake—Regulation and implications for glycaemic control. *Nature Reviews Endocrinology*, 13(3), 133–148. <https://doi.org/10.1038/nrendo.2016.162>
 5. Karelis, A. D., Smith, J. E. W., Passe, D. H., & Péronnet, F. (2010). Carbohydrate Administration and Exercise Performance. *Sports Medicine*, 40(9), 747–763. <https://doi.org/10.2165/11533080-000000000-00000>
 6. da Conceição Chaves de Lemos, M. (2022). Carbohydrate Counting. In F. Bandeira, H. Gharib, L. Griz, & M. Faria (Eds.), *Endocrinology and Diabetes: A Problem Oriented Approach* (pp. 425–431). Springer International Publishing. https://doi.org/10.1007/978-3-030-90684-9_40
 7. Galindo, R. J., & Aleppo, G. (2020). Continuous glucose monitoring: The achievement of 100 years of innovation in diabetes technology. *Diabetes Research and Clinical Practice*, 170, 108502. <https://doi.org/10.1016/j.diabres.2020.108502>
 8. Oliver, N., Gimenez, M., Calhoun, P., Cohen, N., Moscardo, V., Hermanns, N., Freckmann, G., Reddy, M., & Heinemann, L. (2019). Continuous Glucose Monitoring in People With Type 1 Diabetes on Multiple-Dose Injection Therapy: The Relationship Between Glycemic Control and Hypoglycemia. *Diabetes Care*, 43(1), 53–58. <https://doi.org/10.2337/dc19-0977>
 9. Ishihara, K., Uchiyama, N., Kizaki, S., Mori, E., Nonaka, T., & Oneda, H. (2020). Application of Continuous Glucose Monitoring for Assessment of Individual Carbohydrate Requirement during Ultramarathon Race. *Nutrients*, 12(4), Article 4. <https://doi.org/10.3390/nu12041121>
 10. Sengoku, Y., Nakamura, K., Ogata, H., Nabekura, Y., Nagasaka, S., & Tokuyama, K. (2015). Continuous Glucose Monitoring During a 100-km Race: A Case Study in an Elite Ultramarathon Runner. *International Journal of Sports Physiology and Performance*, 10(1), 124–127. <https://doi.org/10.1123/ijsp.2013-0493>
 11. Bowler, A.-L. M., Whitfield, J., Marshall, L., Coffey, V. G., Burke, L. M., & Cox, G. R. (2022). The Use of Continuous Glucose Monitors in Sport: Possible Applications and Considerations. *International Journal of Sport Nutrition and Exercise Metabolism*, 33(2), 121–132. <https://doi.org/10.1123/ijsnem.2022-0139>
 12. American Diabetes Association. (2018). Good to Know: Factors Affecting Blood Glucose. *Clinical Diabetes*, 36(2), 202. <https://doi.org/10.2337/cd18-0012>
 13. Wati, I. D. P., Kusnanik, N. W., Wahjuni, E. S., Samodra, Y. T. J., Gandasari, M. F., & Sofyan, D. (2024). Eat well to the best performance: Calorie intake and eating behavior among athlete: a review. *International Journal of Public Health Science (IJPHS)*, 13(1), 253. <https://doi.org/10.11591/ijphs.v13i1.23336>
 14. Nay, K., Jollet, M., Goustard, B., Baati, N., Vernus, B., Pontones, M., Lefevre-Orfila, L., Bendavid, C., Rué, O., Mariadassou, M., Bonnieu, A., Ollendorff, V., Lepage, P., Derbré, F., & Koechlin-Ramonatxo, C. (2019). Gut bacteria are critical for optimal muscle function: A potential

- link with glucose homeostasis. *American Journal of Physiology-Endocrinology and Metabolism*, 317(1), E158–E171. <https://doi.org/10.1152/ajpendo.00521.2018>
15. Brooks, G. A. (2020). The Precious Few Grams of Glucose During Exercise. *International Journal of Molecular Sciences*, 21(16), Article 16. <https://doi.org/10.3390/ijms21165733>
 16. Bradley, P. S., Carling, C., Gomez Diaz, A., Hood, P., Barnes, C., Ade, J., Boddy, M., Krstrup, P., & Mohr, M. (2013). Match performance and physical capacity of players in the top three competitive standards of English professional soccer. *Human Movement Science*, 32(4), 808–821. <https://doi.org/10.1016/j.humov.2013.06.002>
 17. Alva, S., Bailey, T., Brazg, R., Budiman, E. S., Castorino, K., Christiansen, M. P., Forlenza, G., Kipnes, M., Liljenquist, D. R., & Liu, H. (2022). Accuracy of a 14-Day Factory-Calibrated Continuous Glucose Monitoring System With Advanced Algorithm in Pediatric and Adult Population With Diabetes. *Journal of Diabetes Science and Technology*, 16(1), 70–77. <https://doi.org/10.1177/1932296820958754>
 18. Abbott. (2024). *Home | FreeStyle Libre | Abbott*. <https://www.freestyle.abbott/uk-en/home.html>
 19. Costello, N., Deighton, K., Dyson, J., Mckenna, J., & Jones, B. (2017). Snap-N-Send: A valid and reliable method for assessing the energy intake of elite adolescent athletes. *European Journal of Sport Science*, 17(8), 1044–1055. <https://doi.org/10.1080/17461391.2017.1337815>
 20. Meta Platforms Inc. (2023). *WhatsApp* (Version 24.9.77) [Computer software]. <https://whatsapp.com>
 21. IBM Corp. (2023). *IBM SPSS Statistics for Macintosh* (Version 29.0.2.0) [Computer software]. IBM Corp.
 22. Bangsbo, J., Mohr, M., & Krstrup, P. (2006). Physical and metabolic demands of training and match-play in the elite football player. *Journal of Sports Sciences*, 24(7), 665–674. <https://doi.org/10.1080/02640410500482529>
 23. Ehrhardt, N., & Al Zaghal, E. (2019). Behavior Modification in Prediabetes and Diabetes: Potential Use of Real-Time Continuous Glucose Monitoring. *Journal of Diabetes Science and Technology*, 13(2), 271–275. <https://doi.org/10.1177/1932296818790994>
 24. Chryssanthopoulos, C., Souglis, A., Tsalouhidou, S., Hulton, A. T., Bogdanis, G. C., Petridou, A., Philippou, A., Maridaki, M., & Theos, A. (2024). Dietary Intake of Soccer Players before, during and after an Official Game: Influence of Competition Level and Playing Position. *Nutrients*, 16(3), Article 3. <https://doi.org/10.3390/nu16030337>
 25. Collins, J., Maughan, R. J., Gleeson, M., Billsborough, J., Jeukendrup, A., Morton, J. P., Phillips, S. M., Armstrong, L., Burke, L. M., Close, G. L., Duffield, R., Larson-Meyer, E., Louis, J., Medina, D., Meyer, F., Rollo, I., Sundgot-Borgen, J., Wall, B. T., Boullosa, B., ... McCall, A. (2021). UEFA expert group statement on nutrition in elite football. Current evidence to inform practical recommendations and guide future research. *British Journal of Sports Medicine*, 55(8), 416–416. <https://doi.org/10.1136/bjsports-2019-101961>
 26. Read, D., Cope, E., & Taylor, L. (2024). *Snus Use In English Professional Football*. Loughborough University. <https://www.lborolondon.ac.uk/media/media/london/images/news/2024/snus-use-in-english-professional-football.pdf>
 27. Bergman, B. C., Perreault, L., Hunerdosse, D., Kerege, A., Playdon, M., Samek, A. M., & Eckel, R.

- H. (2012). Novel and Reversible Mechanisms of Smoking-Induced Insulin Resistance in Humans. *Diabetes*, *61*(12), 3156–3166. <https://doi.org/10.2337/db12-0418>
28. Carlsson, S., Andersson, T., Araghi, M., Galanti, R., Lager, A., Lundberg, M., Nilsson, P., Norberg, M., Pedersen, N. L., Trolle-Lagerros, Y., & Magnusson, C. (2017). Smokeless tobacco (snus) is associated with an increased risk of type 2 diabetes: Results from five pooled cohorts. *Journal of Internal Medicine*, *281*(4), 398–406. <https://doi.org/10.1111/joim.12592>
29. Bowler, A.-L. M., Burke, L. M., Coffey, V. G., & Cox, G. R. (2024). Day-to-Day Glycemic Variability Using Continuous Glucose Monitors in Endurance Athletes. *Journal of Diabetes Science and Technology*, 19322968241250355. <https://doi.org/10.1177/19322968241250355>
30. Nutritics. (2019). *Research Edition* (Version v5.09) [Computer software]. <https://www.nutritics.com>
31. Tt1 Products, Inc. (2020). *Supersapiens* (Version 3.1.3) [Computer software]. Abbott. UCI. (2024). *General Organisation of Cycling as a Sport* (E0624). https://assets.ctfassets.net/76117gh5x5an/wQympSG6EWIKq6o6HKw9E/dbf22e49f7cde9ae61df6180c7a90c8e/1-GEN-20240617_E.pdf